

Very viscous drops cannot break up

By Jens Eggers* and Marco A. Fontelos†

*School of Mathematics, University of Bristol, University Walk,
Bristol BS8 1TW, United Kingdom † Departamento de Matemática Aplicada, Universidad
Rey Juan Carlos, C/ Tulipán S/N, 28933 Móstoles, Madrid, Spain.

(Received 2 February 2008)

We consider an axisymmetric, freely suspended fluid drop with surface tension, whose viscosity is so large that both inertia and forcing by an external fluid can be ignored. We show that whatever be the initial condition, pinchoff can never occur.

The breakup of fluid drops has been studied very extensively (Stone (1994), Eggers (1997), Basaran (2002)) owing to its relevance to mixing, printing, and DNA analysis. In most circumstances, breakup of an extended drop occurs almost inevitably owing to the Rayleigh instability (Eggers (1997)), which tends to locally reduce the radius until breakup occurs. So one might think that a sufficiently extended drop, that has separated from a nozzle or has been stretched by an external fluid of comparatively low viscosity, will break up in the same manner. Here we show that this is not the case, but that the drop always retracts to its spherical state of minimum surface energy, if inertial and external forces can be neglected, see figure 1. Stone (1994), p.81, makes some prescient remarks which amount to the same statement, but we are not aware that the impossibility of breakup has ever been shown.

The interior of the drop is thus described by Stokes' equation, subject to a normal stress $\gamma\kappa\mathbf{n}$, where γ is the coefficient of surface tension and $\kappa = 1/h(1+h_z^2)^{1/2} - h_{zz}/(1+h_z^2)^{3/2}$ twice the mean curvature of the interface. If σ is the stress tensor, this can be summarised concisely by

$$\nabla \cdot \sigma = 0 \quad \text{in the drop}, \quad \sigma \mathbf{n} = -\gamma \kappa \mathbf{n} \quad \text{on the surface.} \quad (0.1)$$

Integrating $\nabla \cdot \sigma$ over a volume V bounded by the drop surface and a plane perpendicular to the axis (cf. figure 1), we find from the divergence theorem and from the boundary condition that

$$0 = \int_S \mathbf{n} \sigma ds = \int_{Cr(z)} \mathbf{n} \sigma ds + \int_O \mathbf{n} \sigma ds = \int_{Cr(z)} \mathbf{e}_z \sigma ds - \gamma \int_O \mathbf{n} \kappa ds, \quad (0.2)$$

where O is the surface as shown in figure 1, and $Cr(z)$ is the cross section of the drop at z .

Using $\mathbf{n} = (-h_z \mathbf{e}_z + \mathbf{e}_r)/(1+h_z^2)^{1/2}$, the integral over O can be evaluated as

$$-2\pi\gamma\mathbf{e}_z \int_{z_{end}}^z h h_z \kappa dz = -2\pi\gamma\mathbf{e}_z \int_{z_{end}}^z \left[h/(1+h_z^2)^{1/2} \right]_z dz = -2\pi\gamma\mathbf{e}_z h/(1+h_z^2)^{1/2}, \quad (0.3)$$

since the height $h(z, t)$ goes to zero at the end of the drop. Thus we arrive at the exact relation

$$\int_0^{h(z,t)} \mathbf{e}_z \sigma r dr = -\gamma \mathbf{e}_z h/(1+h_z^2)^{1/2} \quad (0.4)$$




FIGURE 1. A viscous drop of unperturbed radius R is initially extended to a length of $15.5 R$. The full lines show it retracting back into a sphere following Stokes' equation, at $t = n \times 0.646 \eta R / \gamma, n = 0, 1, 2, 3$. The dashed lines are profiles computed on the basis of the lubrication equation (0.6), shown at the same times. Note that the minimum local drop radius is always a monotonically increasing function.

for the total force on the cross section of the drop.

Using the definition of the stress tensor (Landau & Lifshitz 1982), the $\mathbf{e}_z$ -component of (0.4) can be rewritten as

$$\int_0^{h(z)} p(r, z) r dr + 2\eta v_r(h(z), z) = \gamma h / (1 + h_z^2)^{1/2}, \quad (0.5)$$

where $p(r, z)$ is the pressure, η the viscosity of the liquid, and $v_r(r, z)$ the radial component of the velocity.

We want to show that $v_r(h(z), z)$ is always positive at local minima h_{min} of $h(z)$, which would mean that the minimum radius can only increase, so breakup can never occur. We are not able to compute the integral over the pressure in the most general case, so we concentrate on the case that h_{min} is small, as it must be if breakup were to occur. In that case the radial dependence of the pressure is well described by a constant, which is (Eggers (1997)) $p(r, z) \approx \gamma \kappa + 2\eta v_r(h, z)/h$. Using this expression for the pressure, (0.5) turns into

$$v_r(h(z), z) = (\gamma/6\eta) \left(1/(1 + h_z^2)^{1/2} + h h_{zz} / (1 + h_z^2)^{3/2} \right). \quad (0.6)$$

At a local minimum of h h_{zz} is positive, making v_r positive, so h_{min} is *increasing* in time and breakup is impossible.

We have been careful to invoke (0.6) only in the region around h_{min} , where it is sure to hold. In fact, (0.6) is equivalent to the one-dimensional lubrication-type description, which was found to often work surprisingly well throughout a fluid drop or filament (Eggers (1997), Basaran (2002)). Here, we find this observation confirmed, as illustrated in figure 1 by superimposing the lubrication calculation (dashed lines) onto the full numerical calculation. No adjustable parameter was introduced in the comparison. It follows from (0.6) and is illustrated by figure 1, that the minimum radius as given by the lubrication approximation must be monotonically increasing. We suspect that the same holds true for the full Stokes equations, but at present we can't exclude a decreasing h_{min} in cases where h_{min} is not small. It would be interesting to extend our results to the case that an exterior fluid is present. Results presented by Powers et al. (1998) suggest that breakup does not occur as long as the exterior viscosity is sufficiently small.

We gratefully acknowledge funding by the EPSRC.

REFERENCES

- H.A. Stone 1994 Dynamics of drop deformation and breakup in viscous fluids *Annu. Rev. Fluid Mech.* **26**, 65-102.
- Eggers, J. 1997 Nonlinear dynamics and breakup of free surface flows. *Rev. Mod. Phys.* 69, 865-929.
- O. A. Basaran 2002 Small-scale free surface flows with breakup: Drop formation and emerging applications *AIChE J.* **48**, 1842-1848.
- LANDAU L. D. & LIFSCHITZ E. M. 1982 Fluid Mechanics. Oxford
- T. R. Powers et al. 1998 Propagation of a topological transition: The Rayleigh instability *Phys. Fluids* **10**, 1052-1057.